



A service of the National Library of Medicine and the National Institutes of Health

My NCBI [Sign In] [Regis

-> NCBI		up 74	Www.p	ubmed.go	V		[Sign I	n] [Regis		
All Databases	PubMed	Nucleotide	Protein	Genome	Structure	OMIM PN	IC Journals	Bool		
Search PubMed		▼ for 350	1 antibody				Preview Go	Clear		
	Limits			tory Clipbo						
About Entrez Text Version	<ul> <li>Search History will be lost after eight hours of inactivity.</li> <li>Search numbers may not be continuous; all searches are represented.</li> <li>To save search indefinitely, click query # and select Save in My NCBI.</li> <li>To combine searches use #search, e.g., #2 AND #3 or click query # for more</li> </ul>									
Entrez PubMed Overview	Searc	options. ch	Mo	st Recent (	Queries		Time	Result		
Help   FAQ Tutorials	#1	9 Search <b>35</b> 0	C1 antibo	dy			15:33:42	1		
New/Noteworthy E-Utilities	<u>#1</u>	7 Search (br (monoclor		r-amplifie r antibody	,	and	15:31:56	<u>45</u>		
PubMed Services Journals Database MeSH Database	<u># 1</u>	6 Search (ha	rk1) and	(monoclon	al or Ig or	antibody)	15:30:18	<u>0</u>		
	<u>#1</u>	5 Search (au Ig or anti		ted kinase	1) and (mo	noclonal or	15:29:53	<u>0</u>		
Single Citation Matcher Batch Citation	<u>#1</u>	4 Search (au (monoclo	-	related kii r antibody	•	l	15:28:48	<u>4</u>		
Matcher Clinical Queries Special Queries LinkOut My NCBI  Related Resources Order Documents NLM Mobile NLM Catalog NLM Gateway TOXNET Consumer Health	<u>#1</u>	2 Search (se or Ig or a		nine kinas	e 15) and (	monoclonal	15:27:33	<u>29</u>		
	· <u>#1</u>		nase 1) or r (breast-t	(aurora-re	e 15) or (at elated kina blified kina	se 1) or	15:22:51	<u>16</u>		
	<u>#</u>	9 Search (se related ki	rine/threo nase 1) or	(aurora-ro	e 15) or (au elated kina blified kina	se 1) or	15:22:38	898		
Clinical Alerts	<u>#</u>	<u>48</u> Search #7	•	-		-	15:19:40	<u>5</u>		
ClinicalTrials.gov PubMed Central	. ±	7 Search au	rora-A				15:13:11	<u>369</u>		
T UDIVIOU OCITUAL	£	6 Search TA	CI-Ig and	l non-hodg	gkin's		09:01:20	<u>0</u>		
	#	5 Search TA	CI-Fc and	d non-hod	gkin's		09:01:13	<u>0</u>		
	<u>#</u>	3 Search TA	CI-Fc an	d lymphon	na		08:59:04	2		

Clear History

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

#1 Search TACI-Ig or TACI-Fc and non-hodgkin's

08:56:32

10

This is G o o g I e's cache of http://www.ebi.ac.uk/thornton-srv/databases/cgi-bin/pdbsum/GetPage.pl? pdbcode=1ol5 as retrieved on Aug 20, 2007 14:23:36 GMT.

Google's cache is the snapshot that we took of the page as we crawled the web.

The page may have changed since that time. Click here for the current page without highlighting.

This cached page may reference images which are no longer available. Click here for the cached text only.

To link to or bookmark this page, use the following url: http://www.google.com/search? q=cache:dpdyHouwH6YJ:www.ebi.ac.uk/thornton-srv/databases/cgi-bin/pdbsum/GetPage.pl%3Fpdbcode% 3D1ol5+aurora-A+and+synonym&hl=en&ct=clnk&cd=4&gl=us

Google is neither affiliated with the authors of this page nor responsible for its content.

These search terms have been highlighted: aurora a synonym



All Databases

Enter Text Here



Go to PDB code: 10



Top page 🔭 🕲 Protein 🕲 Ligands 🕲 Prot-prot 🔮 Clef

Complex(kinase/cell division protein)

PDB id: 1ol5

Name: Complex(kinase/cell division protein)

Title: Structure of aurora-a 122-403, phosphorylated on thr287,

thr288 and bound to tpx2 1-43

Structure: Serine/threonine kinase 6. Synonym: aurora-a,

serine/threonine kinase 15, aurora/ipi1-related kinase 1, aurora-related kinase 1, hark1, breast-tumor-amplified kinase. Chain: a. Fragment: catalytic domain residues 122 403. Engineered: yes. Other\_details: phosphorylated on

thr287, thr288. Restricted expression proliferation

associated

Source: Homo sapiens. Human. Expressed in: escherichia coli.

Biological unit: Dimer (from PDB file)

UniProt: Chain A: O14965 (STK6\_HUMAN)

Struc: Sea

403 a

مزارز درزز اسرزز سرززز ازززز درزار

66 a

# Contents

#### Description

Header details

Header records

Biological unit = asymmetric unit,

as shown

(as defined in PDB file)

References

**PROCHECK** 

#### **Protein chains**

266 a.a. \*

<u>30 a.a.</u> \*

Ligands

Struc	200 a
Chain B: Q9ULW0 (TPX2_HUMAN) [Pfan	<u>n]</u> .
Seq:	2
مرزز المستعدد	
Struc: Sea:	
5/8/2//////////////////////////////////	Z
Struc:	••
Seq: 1935///////	747 :
55000000000000000000000000000000000000	
Charles	20 -



\* PDB and UniProt seqs differ at 2 residue positions (black crosses)

Metal ions ■ MG ×3

Waters ×144

\* Residue conservation analysis

Tools

Image Generation

AstexViewer™@MSD-EBI

Run PROCHECK

Clefts Calculation

Enzyme class: Chain A: E.C.2.7.1.37 [IntEnz] [ExPASy] [KEGG]

[BRENDA]

Reaction: ATP + a protein = ADP + a phosphoprotein (see diagram

below)

Function: (see GO annotation below)

Resolution: 2.50Å R-factor: 0.194 R-free: 0.252

Authors: R.Bayliss, E.Conti

Key ref: R.Bayliss et al. (2003). Structural basis of Aurora-A Cell activation by TPX2 at the mitotic spindle.. *Mol Cell*, 12, 85

862. [PubMed id: 14580337] [DOI: 10.1016/S1097-2765(0

00392-7] **●**Date: 06-Aug-03

Release date: 30-Oct-03

Related entries: 1muo crystal structure of aurora-2, an oncogenic serine-

threonine kinase

1016 structure of unphosphorylated d274n mutant of auroi

а

1017 structure of human aurora-a 122-403 phosphorylated

on thr287, thr288



### Gene Ontology (GO) functional annotation

Biological process protein amino acid phosphorylation 1 term(s)
Biochemical function protein kinase activity 3 term(s)

For full annotation, click on icon

Enzyme reaction for E.C.2.7.1.37

Molecule diagrams generated from .mol files obtained from the KEGG ftp site.

Key re1 Mol Cell 12:85

DOI no: 10.1016/S1097-2765(03)00392-7

PubMed id: 14580337

# Structural basis of Aurora-A activation by TPX2 at the mitotic spindle R.Bayliss, T.Sardon, I.Vernos, E.Conti.

### **ABSTRACT**

Aurora-A is an oncogenic kinase essential for mitotic spindle assembly. It is activated by phosphorylation and by the microtubule-associated protein TPX2, which also localizes the kinase to spindle microtubules. We have uncovered the

ephosphorylation. In the absence of TPX2, the Aurora-A activation segment is in an inactive conformation, with the cru phosphothreonine exposed and accessible for deactivation. Binding of TPX2 triggers no global conformational changes kinase but pulls on the activation segment, swinging the phosphothreonine into a buried position and locking the active conformation. The recognition between Aurora-A and TPX2 resembles that between the cAPK catalytic core and its flan regions, suggesting this molecular mechanism may be a recurring theme in kinase regulation.



### Selected figure(s)





Figure 3. Figure 3. Structure of Aurora-A Bound to TPX2(A) View of the complex between the catalytic domain of human Aurora (Aurora N, yellow) and the N-terminal domain of TPX2 shown in typical kinase orientation. An upstream stretch of TPX2 (red) binds at the N-terminal lobe of Aurora-A, and a downstream stretch (pink) binds between the two lobes. A dotted line in pink marks the approximate path of the linker connecting the two TPX2 stretches (disordered and not modeled).(B) View of the complex after a 180° rotation about the vertical axis in respect to view in (A) shows more clearly the two stretches of TPX2 binding to Aurora-A.(C) The upstream stretch of TPX2 (red, residues 7-21^TPX) binds at a hydrophobic surface groove present in the N-terminal lobe of the kinase (gray cartoon, yellow side chains). Details of the extensive interactions are shown in the same orientation as in (B). Aurora-A residues are labeled in black, and TPX2 residue labels are color coded as the structure.(D) The downstream helical stretch of TPX2 (pink, residues 30-43<sup>Δ</sup>TPX) binds Aurora-A near helix αC and the activation segment, close to but not directly in contact with phospho-Thr288^AUR

(green). Details of interactions are shown in the same orientation as in

(B) and (C).




Figure 5. Figure 5. TPX2-Aurora-A Intermolecular Interactions Resemble cAF Interactions(A and B) Transparent surfaces representing the conser cores of (A) Aurora-A and (B) cAPK show similar surface grooves in lobe (between helix  $\alpha C$  and the  $\beta$  sheet, gray cartoon) and a similar the two lobes (formed by the activation segment and helix αC, gray portions of TPX2 binding to Aurora-A are shown in red and pink (A), C-terminal extensions to the cAPK catalytic core are shown in light I D) Schematic diagram of the intermolecular interactions between At TPX2 (pink and red) and of the cAPK intramolecular interactions (lig that their mode of recognition at the atomic level is rather similar. Th interactions of Tyr8^TPX, Tyr10^TPX, Trp34^TPX, and Phe35^TPX recapitulated by Phe347^cAPK, Phe350^cAPK, Trp30^cAPK, and F

The above figures are reprinted by permission from Cell Press: Mol Cell (2003, 12, 851-862) copyright 2003. Figures were selected by an automated process.

Terms of Use : EBI Funding : Contact EBI : © European Bioinformatics Institute 2006-2007. EBI is an Outstation of the European Mole





A service of the National Library of Medicine and the National Institutes of Health

My NCBI [<u>Sign In]</u> [Regis<sup>.</sup>

www.pubmed.gov All Databases PubMed Nucleotide Protein Genome Structure OMIM **PMC** Journals Search PubMed Preview: Go Clear for Preview/Index History Clipboard • Search History will be lost after eight hours of inactivity. About Entrez • Search numbers may not be continuous; all searches are represented. • To save search indefinitely, click query # and select Save in My NCBI. **Text Version** • To combine searches use #search, e.g., #2 AND #3 or click query # for more options. Entrez PubMed **Most Recent Queries** Search Time Result Overview Help | FAQ #17 Search (breast-tumor-amplified kinase) and 15:53:53 45 Tutorials (monoclonal or Ig or antibody) New/Noteworthy E-Utilities #12 Search (serine/threonine kinase 15) and (monoclonal 29 or Ig or antibody) PubMed Services #14 Search (aurora/ipl1-related kinase 1) and 15:42:18 4 Journals Database (monoclonal or Ig or antibody) MeSH Database Single Citation #19 Search 35C1 antibody 15:33:42 1 Matcher **Batch Citation**  $\frac{\#16}{}$  Search (hark1) and (monoclonal or Ig or antibody) 15:30:18 0 Matcher #15 Search (aurora-related kinase 1) and (monoclonal or 15:29:53 0 Clinical Queries Special Queries Ig or antibody) LinkOut #10 Search (serine/threonine kinase 15) or (aurora/ipl1-15:22:51 16 My NCBI related kinase 1) or (aurora-related kinase 1) or (hark1) or (breast-tumor-amplified kinase) and Related Resources **Order Documents** monoclonal **NLM Mobile** #9 Search (serine/threonine kinase 15) or (aurora/ipl1-15:22:38 898 **NLM Catalog NLM Gateway** related kinase 1) or (aurora-related kinase 1) or TOXNET (hark1) or (breast-tumor-amplified kinase) Consumer Health Clinical Alerts #8 Search #7 and monoclonal 15:19:40 <u>5</u> ClinicalTrials.gov #7 Search aurora-A 15:13:11 369 PubMed Central #6 Search TACI-Ig and non-hodgkin's 09:01:20 0 0 #5 Search TACI-Fc and non-hodgkin's 09:01:13 2 #3 Search TACI-Fc and lymphoma 08:59:04

Clear History

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

#1 Search TACI-Ig or TACI-Fc and non-hodgkin's

08:56:32

10

## **WEST Search History**

Hide Items Restore Clear Cancel

DATE: Monday, September 10, 2007

Hide?	<u>Set</u> Name	Query	Hit Count
	DB=P	GPB, USPT, USOC, EPAB, JPAB, DWPI; PLUR=YES; OP=ADJ	
	L5	L3 and aurora-A kinase	. 0
	L4	L3 and aurora-A	1
	L3	L2 and (monoclonal or Ig or antibody.)	38
	L2	(serine/threonine kinase 15) or (aurora/ipl1-related kinase 1) or (aurora-related kinase 1) or (hark1) or (breast-tumor-amplified kinase)	39
	DB=P	GPB, USPT; PLUR=YES; OP=ADJ	
	L1	(serine/threonine kinase 15) or (aurora/ipl1-related kinase 1) or (aurora-related kinase 1) or (hark1) or (breast-tumor-amplified kinase)	39

**END OF SEARCH HISTORY**